

Lesson 10: A Critical Look at Proportional Relationships

Classwork

Example 1

Paul walks 2 miles in 25 minutes. How many miles can Paul walk in 137.5 minutes?

Time (in minutes)	Distance (in miles)
25	2

- b. What fraction represents constant speed, C , if it takes her x number of minutes to get halfway to her friend's house?
- c. Write and solve a proportion using the fractions from parts (a) and (b) to determine how many minutes it takes her to get to the halfway point.
- d. Write a two-variable equation to represent how many miles Stefanie can drive over any time interval.
3. The equation that represents how many miles, y , Dave travels after x hours is $y = 50x + 15$. Use the equation to complete the table below.

x (hours)	Linear Equation: $y = 50x + 15$	y (miles)
1	$y = 50(1) + 15$	65
2		
3		
3.5		
4.1		

Lesson Summary

Average speed is found by taking the total distance traveled in a given time interval, divided by the time interval.

If y is the total distance traveled in a given time interval x , then $\frac{y}{x}$ is the average speed.

If we assume the same average speed over any time interval, then we have constant speed, which can then be used to express a linear equation in two variables relating distance and time.

If $\frac{y}{x} = C$, where C is a constant, then you have constant speed.

Problem Set

- Eman walks from the store to her friend's house, 2 miles away. It takes her 35 minutes.
 - What fraction represents her constant speed, C ?
 - Write the fraction that represents her constant speed, C , if she walks y miles in 10 minutes.
 - Write and solve a proportion using the fractions from parts (a) and (b) to determine how many miles she walks after 10 minutes. Round your answer to the hundredths place.
 - Write a two-variable equation to represent how many miles Eman can walk over any time interval.
- Erika drives from school to soccer practice 1.3 miles away. It takes her 7 minutes.
 - What fraction represents her constant speed, C ?
 - What fraction represents her constant speed, C , if it takes her x minutes to drive exactly 1 mile?
 - Write and solve a proportion using the fractions from parts (a) and (b) to determine how much time it takes her to drive exactly 1 mile. Round your answer to the tenths place.
 - Write a two-variable equation to represent how many miles Erika can drive over any time interval.
- Darla drives at a constant speed of 45 miles per hour.
 - If she drives for y miles and it takes her x hours, write the two-variable equation to represent the number of miles Darla can drive in x hours.
 - Darla plans to drive to the market 14 miles from her house, then to the post office 3 miles from the market, and then return home, which is 15 miles from the post office. Assuming she drives at a constant speed the entire time, how long will it take her to run her errands and get back home? Round your answer to the hundredths place.
- Aaron walks from his sister's house to his cousin's house, a distance of 4 miles, in 80 minutes. How far does he walk in 30 minutes?

5. Carlos walks 4 miles every night for exercise. It takes him exactly 63 minutes to finish his walk.
- Assuming he walks at a constant rate, write an equation that represents how many miles, y , Carlos can walk in x minutes.
 - Use your equation from part (a) to complete the table below. Use a calculator, and round all values to the hundredths place.

x (minutes)	Linear Equation:	y (miles)
15		
30		
40		
60		
75		